

# SYNTHESIS AND SOME REACTIONS OF 11-AZACYCLOHEPT[*a*]AZULEN-3(3*H*)-ONES AND EVALUATION OF THEIR CYTOTOXIC ACTIVITY AGAINST HELA S3 CELLS

Tomoyuki Ariyoshi,<sup>1</sup> Kengo Yoshinaga,<sup>2</sup> Kazuya Koizumi,<sup>1</sup> Hiroyuki Fujii,<sup>3</sup> Reiko Ikeda,<sup>4</sup> Takeo Konakahara,<sup>4</sup> and Noritaka Abe<sup>1\*</sup>

<sup>1</sup>Applied Molecular Bioscience, Graduate School of Medicine, and Department of Biology and Chemistry, Faculty of Science, Yamaguchi University, Yoshida, Yamaguchi 753-8512, Japan

<sup>2</sup>Department of Biology and Chemistry, Faculty of Science, Yamaguchi University, Yamaguchi 753-8512, Japan

<sup>3</sup>Science Research Center, Yamaguchi University, Yamaguchi 753-8512, Japan

<sup>4</sup>Department of Pure and Applied Chemistry, Faculty of Science and Technology, Tokyo University of Science, Noda, Chiba 278-8510, Japan

**Abstract** – Aldol condensation of 2,3-diformyl-1-azaazulene with acetone, methoxyacetone, and benzylacetone in aq. NaOH solution gave corresponding 11-azacyclohept[*a*]azulen-3(3*H*)-one (**2a**), 2-methoxy-derivative (**2b**) and 2-benzyl-derivative (**2d**) in good yields. Hydrolysis of **2b** using hot HBr gave 2-hydroxy-11-azacyclohept[*a*]azulen-3(3*H*)-one (**2f**). Acetylation of **2f** gave 2-acetoxy-11-azacyclohept[*a*]azulen-3(3*H*)-one (**2g**). Reaction of the compounds (**2a**, **2b**, **2f**) with diethyl malonate in Ac<sub>2</sub>O under heating gave 1,9(11*bH*)-11*b*-azaazuleno[1,2,3-*cd*]azulenedione derivatives (**4a**, **4b**, **4f**). Compounds (**2b**, **2f**) showed weak cytotoxic activity against HeLa S3 cells.

## INTRODUCTION

Tricyclic and tetracyclic systems of containing azulene nucleus such as azulenotropylium ions, azulenotropones and azulenoazulenes are attracted attentions from the view of aromaticity and bioactivity.<sup>1-3</sup> Several azulenotropones and azulenoazulenes have been synthesized notably by K. Takase,<sup>4-6</sup> C. Jutz,<sup>7</sup> T. Toda,<sup>8</sup> Z. Yoshida,<sup>9</sup> and S. Kuroda,<sup>10</sup> and M. Nitta.<sup>11</sup> Although many researches



2-methoxy-11-azacyclohept[*a*]azulen-3(3*H*)-one (**2b**) in 63% yield, and 4-methoxy-11-azacyclohept[*a*]azulen-3(3*H*)-one<sup>13</sup> (**2c**) was not obtained. Melting point of obtained compound (mp 214-215 °C) differed from that of the reported **2c** (mp 209.5-211.5 °C)<sup>13</sup>. The electronic spectra of **2b** and **2c**<sup>13</sup> were distinctly different as shown in Fig. 2. In the <sup>1</sup>H NMR spectrum of **2b**, methoxy proton was seen at  $\delta$  4.11, and a singlet signal assigned to H-1 proton was seen at  $\delta$  7.68 and AB-doublet protons were seen at  $\delta$  7.13 (1H, d, *J* 12.2, H-4) 8.17 (1H, d, *J* 12.2, H-5), except for the seven-membered ring protons of the azaazulene moiety. From the results, we assigned the structure.




Figure 1. Calculated bond lengths and the optimized structure of **2a**.

When **1** was treated with benzylacetone for 10 min at rt, **2d** (34%) and **3** (22%) were obtained. In the <sup>1</sup>H NMR spectrum of **3**, two methine protons were observed at  $\delta$  3.56 (1H, dddd, *J* 8.1, 7.7, 6.0, and 1.2, H-2), 5.21 (1H, d, *J* 6.0, H-1) and benzyl protons were seen at  $\delta$  2.60 (1H, dd, *J* 13.8 and 7.7) and 2.77 (1H, dd, *J* 13.8 and 8.1), except for the seven-membered ring protons of the azaazulene moiety. In the <sup>13</sup>C NMR spectrum of **3**, two methine carbons were observed at  $\delta$  59.1 and 68.8 together with benzyl carbon at  $\delta$  34.3. In the IR spectrum of **3**, characteristic peaks were observed at 3444 (OH), 1623 (C=O), and 1603 cm<sup>-1</sup> (C=C). From the results, we assigned the structure. Prolonged treatment of **1** with benzylacetone (for 2h at rt) gave **2d** in 74% yield and none of **3**. It is considered that intermediate (**3**) was obtained at first, then successive dehydration afforded **2d**.

Similar treatment of **1** with diethyl 1,3-acetonedicarboxylate gave **2e** in 13% yield. For improvement of the yield, **1** was treated under azeotropic conditions in the presence of NHEt<sub>2</sub> in benzene under reflux for 4 h, but **2e** was obtained only in 5% yield.

Hydrolysis of **2b** was achieved by treatment with 48% HBr under reflux for 3 h and 2-hydroxy-11-azacyclohept[*a*]azulen-3(3*H*)-one (**2f**) was obtained in 88% yield. Compound **2f** was acetylated with Ac<sub>2</sub>O in the presence of pyridine for 1 h at rt, and 2-acetoxy-11-azacyclohept[*a*]azulen-3(3*H*)-one (**2g**) was obtained in 61% yield, where the isomeric acetate could not be isolated.





Figure 2. Electronic spectra of **2b** and **2c**.

It is well known that 8,8-dicyanoheptafulvene was obtained by the condensation of tropone with malononitrile in refluxing Ac<sub>2</sub>O.<sup>19</sup> Therefore we examined the reaction of **2a** with malononitrile, but distinct product was not isolated. When **2a** was treated with diethyl malonate in refluxing Ac<sub>2</sub>O for 1.5 h gave 11b-azaazuleno[1,2,3-*cd*]azulene-1,9(11*bH*)-dione derivative (**4a**) in 24 % yield. In the <sup>1</sup>H NMR spectrum of **4a**, one ethyl ester signals were observed at δ 1.45 (*J* 7.1, Me) and 4.43 (*J* 7.1, OCH<sub>2</sub>) and four proton signals owing to seven-membered ring of azaazulene moiety were seen at δ 7.51 (dd, *J* 10.4 and 10.3, H-5), 7.81 (dd, *J* 11.0 and 10.3, H-4), 8.04 (d, *J* 10.4, H-6), and 8.71 (d, *J* 11.0, H-3), and tropone moiety at δ 7.07 (dd, *J* 12.2 and 2.5, H-8), 7.22 (dd, *J* 12.4 and 2.5, H-10), 7.73 (dd, *J* 12.2, H-7), and 8.48 (d, *J* 12.4, H-11). In the IR spectrum of **4a**, three carbonyl signals were seen at 1738, 1674, and 1619 cm<sup>-1</sup>, owing to ester, γ-lactim, and tropone carbonyls, respectively. From the results, we

assigned the structure. Michael addition of the enol form of diethyl malonate to the C-10 of **2a** would generate intermediate **A**, and successive cyclization and dehydrogenation furnished to **4a**.

Similar treatment of **2b** and **2d** with diethyl malonate in refluxing  $\text{Ac}_2\text{O}$  gave **4b** (39%) and **4f** (25%), respectively.

As **2a** was not reacted at the carbonyl of tropone moiety as above, we examined the reaction of **2a** with 2,4-dinitrophenyl hydrazine to substantiate the reactivity. But we could not obtain distinct products, and failed to elicit the reactivity of the carbonyl group.



### Evaluation of cytotoxic activity

Compounds (**2b** and **2f**) were evaluated for their cytotoxic activity against HeLa S3 cells. The  $\text{IC}_{50}$  values [ $\mu\text{M}$ ] were 41 and 19 for **2b** and **2f**, respectively. The results revealed that the compounds (**2b** and **2f**) showed weak activity against HeLa S3 cells.

### EXPERIMENTAL

Mps are measured using a Yanagimoto micro-melting apparatus and uncorrected.  $^1\text{H}$  NMR spectra (including HH-COSY and CH-COSY NMR) were recorded on a Bruker AVANCE 400S (400 MHz) and  $^{13}\text{C}$  NMR spectra were recorded on a Bruker AVANCE 400S (100.6 MHz) using  $\text{CDCl}_3$  as a solvent with tetramethylsilane as an internal standard unless otherwise stated;  $J$  values are recorded in Hz. IR spectra were recorded for KBr pellets on a Nicolet FT-IR AVTAR 370DTGS unless otherwise stated. Electronic spectra were recorded with JASCO V-570 spectrophotometer. Elemental analyses were taken with a Perkin Elmer 2400II. Kieselgel 60 and activated alumina C300 were used for column chromatography and Kieselgel 60G was used for thin-layer chromatography.

### Synthesis of 11-azacyclohept[*a*]azulen-3(3*H*)-one (**2a**)

A solution of 2,3-diformyl-1-azaazulene (**1**) (0.0988 g, 0.534 mmol) and  $\text{Me}_2\text{CO}$  (0.06 mL, 0.82 mmol) in 0.11 M NaOH solution (100 mL, 1.1 mmol) was stirred for 10 min at room temperature. The mixture was extracted with  $\text{CH}_2\text{Cl}_2$ . The extract was dried over  $\text{Na}_2\text{SO}_4$ , and evaporated. The residue was

chromatographed with EtOAc to give **2a** (0.0531 g, 48%) as purple powders.

**2a:** Purple micro needles (from EtOH), mp 195 °C (decomp.);  $\delta_{\text{H}}$  6.95 (1H, dd,  $J$  12.1 and 2.4, H-4), 7.25 (1H, d,  $J$  12.4 and 2.4, H-2), 7.96 (1H, ddd,  $J$  9.8, 9.6, and 0.4, H-7), 8.03 (1H, dd,  $J$  10.0 and 9.8, H-9), 8.07 (1H, d,  $J$  12.1, H-5), 8.08 (1H, d,  $J$  12.4, H-1), 8.15 (1H, t,  $J$  9.8, H-8), 8.88 (1H, dd,  $J$  10.0 and 0.4, H-10), and 8.93 (1H, d,  $J$  9.6, H-6);  $\delta_{\text{H}}$ (TFA-*d*) 7.71 (1H, dd,  $J$  12.0 and 2.0, H-4), 7.92 (1H, d,  $J$  12.1 and 2.0, H-2), 8.47 (1H, d, 12.1, H-1), 8.83 (1H, dd,  $J$  10.0 and 9.9, H-7), 8.86 (1H, dd,  $J$  10.1 and 9.8, H-9), 8.88 (1H, d,  $J$  12.0, H-5), 8.98 (1H, dd,  $J$  10.0 and 9.8, H-8), 9.39 (1H, d,  $J$  10.1, H-10), and 9.87 (1H, d,  $J$  9.9, H-6);  $\delta_{\text{C}}$  107.5, 120.0, 130.4, 131.9, 132.5, 135.8, 138.7, 139.9, 134.0, 141.0, 141.4, 143.6, 149.7, and 184.9;  $\nu_{\text{max}}$  /  $\text{cm}^{-1}$  1592 (C=O);  $\lambda_{\text{max}}$ (CHCl<sub>3</sub>) nm (log  $\epsilon$ ) 286 (4.11), 328 (4.52), 342 (4.66), 360 (3.97, sh), 385 (3.86), 406 (3.42), 500 (2.82, sh), 541 (3.03, sh), 568 (3.07), and 616 (2.84, sh);  $\lambda_{\text{max}}$ (MeOH) nm (log  $\epsilon$ ) 286 (4.16), 328 (4.60), 342 (4.72), 358 (4.06, sh), 384 (3.93), 406 (3.49), 541 (3.15, sh), 566 (3.17), and 607 (2.97, sh). *Anal.* Calcd for C<sub>14</sub>H<sub>9</sub>NO·H<sub>2</sub>O: C, 74.65; H, 4.92; N, 6.22. Found: C, 74.44; H, 4.98; N, 6.19.

### Synthesis of 2-methoxy-11-azacyclohept[*a*]azulen-3(3*H*)-one (2b)

A solution of **1** (0.0981 g, 0.530 mmol) and methoxyacetone (0.07 mL, 0.76 mmol) in 0.055 M NaOH solution (200 mL, 1.1 mmol) was stirred for 10 min at rt. The mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The extract was dried over Na<sub>2</sub>SO<sub>4</sub>, and evaporated. The residue was chromatographed with EtOAc to give **2b** (0.0795 g, 63%) as reddish purple leaflets.

**2b:** Reddish purple leaflets (from hexane-CH<sub>2</sub>Cl<sub>2</sub>), mp 214-215 °C;  $\delta_{\text{H}}$  4.11 (3H, s, OMe), 7.13 (1H, d,  $J$  12.2, H-4), 7.68 (1H, s, H-1), 7.95 (1H, td,  $J$  9.4 and 1.2, H-7), 7.99 (1H, ddd,  $J$  9.6, 9.4, and 1.2, H-9), 8.10 (1H, ddd,  $J$  9.6, 9.4, and 1.4, H-8), 8.17 (1H, d,  $J$  12.2, H-5), 8.77 (1H, dd,  $J$  9.6 and 1.4, H-10), and 8.86 (1H, d,  $J$  9.4, H-6);  $\delta_{\text{H}}$ (TFA-*d*) 4.35 (3H, s, OMe), 7.92 (1H, d,  $J$  11.9, H-4), 8.13 (1H, s, H-1), 8.81-8.96 (3H, m, H-7, 8, and 9), 9.06 (1H, d,  $J$  11.9, H-5), 9.36 (1H, d,  $J$  9.9, H-10), and 9.82 (1H, d,  $J$  9.8, H-6);  $\delta_{\text{C}}$  56.7, 110.1, 121.9, 128.2, 129.4, 130.9, 132.9, 133.1, 137.1, 139.0, 144.9, 160.0, 161.3, 162.4, and 181.7;  $\nu_{\text{max}}$  /  $\text{cm}^{-1}$  1614 (C=O);  $\lambda_{\text{max}}$ (CHCl<sub>3</sub>) nm (log  $\epsilon$ ) 259 (3.99), 284 (4.19, sh), 296 (4.22), 322 (4.34, sh), 343 (4.64), 358 (4.43, sh), 382 (4.05, sh), 409 (4.00), 528 (3.14, sh), 555 (3.19), and 592 (2.96, sh);  $\lambda_{\text{max}}$ (MeOH) nm (log  $\epsilon$ ) 258 (3.88), 285 (4.19, sh), 294 (4.22), 318 (4.29, sh), 341 (4.63), 376 (4.07, sh), 400 (3.98), and 535 (3.12). *Anal.* Calcd for C<sub>15</sub>H<sub>11</sub>NO<sub>2</sub>·2H<sub>2</sub>O: C, 70.02; H, 5.88; N, 5.44. Found: C, 70.12; H, 5.88; N, 5.34.

### Reaction of **1** with benzylacetone

a) A solution of **1** (0.093 g, 0.502 mmol) and benzylacetone (0.113 g, 0.762 mmol) in 0.048 M NaOH solution (100 mL, 4.80 mmol) was stirred for 10 min at rt. The mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub>.

The extract was dried over Na<sub>2</sub>SO<sub>4</sub>, and evaporated. The residue was chromatographed with EtOAc to give **2d** (0.0471 g, 34%) and **3** (0.0334 g, 22%), successively.

b) A solution of **1** (0.0370 g, 0.200 mmol) and benzylacetone (0.113 g, 0.762 mmol) in 0.0445 M NaOH solution (100 mL, 4.45 mmol) was stirred for 2 h at rt. The mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The extract was dried over Na<sub>2</sub>SO<sub>4</sub>, and evaporated. The residue was chromatographed with EtOAc to give **2d** (0.0442 g, 74%).

**2d**: violet powders (from hexane-CH<sub>2</sub>Cl<sub>2</sub>), mp 208-210 °C;  $\delta_{\text{H}}$  4.14 (2H, s, CH<sub>2</sub>), 7.01 (1H, d, *J* 12.2, H-4), 7.23 (1H, tt, *J* 7.2 and 2.2, H-*p*-Ph), 7.31 (2H, dd, *J* 7.7 and 7.2, H-*m*-Ph), 7.37 (2H, dd, *J* 7.7 and 2.2, H-*o*-Ph), 7.91 (1H, dd, *J* 9.8 and 9.5, H-7), 7.97 (1H, dd, *J* 9.9 and 9.6, H-9), 8.01 (1H, d, *J* 12.2, H-5), 8.08 (1H, dd, *J* 9.9 and 9.8, H-8), 8.12 (1H, s, H-1), 8.79 (1H, d, *J* 9.6, H-10), and 8.86 (1H, d, *J* 9.5, H-6);  $\delta_{\text{C}}$  40.9, 124.0, 126.5, 128.3, 128.6, 129.6, 130.5, 131.1, 132.3, 124.5, 134.6, 138.6, 139.2, 140.1, 144.5, 152.1, 159.1, 161.8, and 187.3;  $\nu_{\text{max}}$  / cm<sup>-1</sup> 1597 (C=O). *Anal.* Calcd for C<sub>21</sub>H<sub>15</sub>NO·H<sub>2</sub>O: C, 79.98; H, 5.43; N, 4.44. Found: C, 79.66; H, 5.32; N, 4.64.

**3**: Red powders (from hexane-CH<sub>2</sub>Cl<sub>2</sub>), mp 202 °C (decomp.);  $\delta_{\text{H}}$  2.60 (1H, dd, *J* 13.8 and 7.7, CH<sub>2</sub>), 2.77 (1H, dd, *J* 13.8 and 8.1, CH<sub>2</sub>), 3.56 (1H, dddd, *J* 8.1, 7.7, 6.0, and 1.2, H-2), 5.21 (1H, d, *J* 6.0, H-1), 6.24 (1H, d, *J* 12.3 and 1.2, H-4), 6.94 (2H, dt, *J* 7.5 and 1.5, H-*o*-Ph), 7.11 (1H, tt, *J* 7.0 and 1.5, H-*p*-Ph), 7.16 (2H, dd, *J* 7.5 and 7.0, H-*m*-Ph), 7.56 (1H, d, *J* 12.3, H-5), 7.76 (1H, dd, *J* 10.0 and 9.7, H-7), 7.80 (1H, dd, *J* 9.8 and 9.7, H-9), 7.97 (1H, ddd, *J* 9.8, 9.7, and 0.8, H-8), 8.57 (1H, d, *J* 10.0, H-6), and 8.63 (1H, dd, *J* 9.5 and 0.8, H-10), (OH was not observed);  $\delta_{\text{C}}$  34.3, 59.1, 68.8, 119.5, 125.9, 126.3, 128.3, 128.9, 130.5, 131.6, 133.8, 137.6, 138.3, 139.7, 144.5, 157.0, 169.9, and 196.9;  $\nu_{\text{max}}$  / cm<sup>-1</sup> 3444 (OH), 1623 (C=O), and 1603 (C=C). *Anal.* Calcd for C<sub>21</sub>H<sub>17</sub>NO<sub>2</sub>: C, 79.98; H, 5.43; N, 4.44. Found: C, 80.06; H, 5.52; N, 4.28.

### Reaction of **1** with diethyl acetone-1,3-dicarboxylate

a) A solution of **1** (0.0951 g, 0.514 mmol) and diethyl acetone-1,3-dicarboxylate (0.20 mL, 1.20 mmol) in 0.25 M NaOH solution (50 mL, 1.25 mmol) was stirred for 10 min at rt. The mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The extract was dried over Na<sub>2</sub>SO<sub>4</sub>, and evaporated. The residue was chromatographed with EtOAc to give **2e** (0.0215 g, 13%) as violet powders.

b) To the solution of diethyl acetone-1,3-dicarboxylate (0.20 mL, 1.20 mmol) and NHEt<sub>2</sub> (0.15 mL) in dry benzene (50 mL) was added benzene solution (20 mL) of **1** (0.1149 g, 0.625 mmol), and the mixture was refluxed for 4 h using Dean-Stark trap. The mixture was evaporated and the residue was chromatographed with EtOAc to give **2e** (0.0015 g, 5%)

**2e**: Violet powders (from AcOEt), mp 148-149 °C;  $\delta_{\text{H}}$  1.41 (3H, t, *J* 7.0, Me), 1.42 (3H, t, *J* 7, Me), 4.43 (4H, q, *J* 7.0, CH<sub>2</sub>), 8.05 (1H, dd, *J* 10.1 and 9.9, H-9), 8.10 (1H, dd, *J* 9.8 and 9.5, H-7), 8.23 (1H, dd, *J*

9.9 and 9.5, H-8), 8.55 (1H, s, H-1), 8.77 (1H, s, H-5), 8.93 (1H, d, *J* 10.1, H-10), and 9.05 (1H, d, *J* 9.5, H-6);  $\delta_{\text{C}}$  14.1, 14.2, 61.9, 62.1, 122.9, 130.9, 131.5, 131.6, 133.3, 134.2, 136.3, 140.3, 141.1, 141.8, 145.9, 159.8, 160.8, 166.2, 167.1, and 184.2;  $\nu_{\text{max}} / \text{cm}^{-1}$  1736, 1695, and 1626 (C=O);  $\lambda_{\text{max}}(\text{CHCl}_3)$  nm (log  $\epsilon$ ) 296 (4.32, sh), 334 (4.59), 341 (4.58), 376 (4.12, sh), 523 (3.00, sh), 553 (3.04), and 592 (2.84, sh). *Anal.* Calcd for  $\text{C}_{20}\text{H}_{17}\text{NO}_5$ : C, 68.37; H, 4.88; N, 3.99. Found: C, 68.57; H, 4.93; N, 3.75.

### Synthesis of 2-hydroxy-11-azacyclohept[*a*]azulen-3(3*H*)-one (2f)

A mixture of **2b** (0.0317 g, 0.134 mmol) and 48% HBr (10 mL) was stirred for 3 h under heating at 160 °C. The mixture was neutralized with  $\text{Na}_2\text{CO}_3$  and extracted with  $\text{CHCl}_3$ . The extract was dried over  $\text{Na}_2\text{SO}_4$ , and evaporated. The compound **2f** (0.0263 g, 88%) was obtained as purple powders.

**2f**: Purple scales (from EtOH-EtOAc), mp 178-180 °C;  $\delta_{\text{H}}$  5.11 (1H, br, OH), 7.31 (1H, d, *J* 12.0, H-4), 7.97 (1H, ddd, *J* 9.6, 9.5 and 1.4, H-7), 8.04 (1H, ddd, *J* 9.6, 9.3, and 1.4, H-9), 8.07 (1H, s, H-1), 8.08 (1H, dd, *J* 9.6 and 9.5, H-8), 8.43 (1H, d, *J* 12.0, H-5), 8.82 (1H, dd, *J* 9.3 and 1.8, H-10), and 8.88 (1H, d, *J* 9.5, H-6);  $\delta_{\text{C}}$  11.9, 122.8, 123.8, 125.0, 131.9, 132.7, 133.4, 135.2, 137.4, 139.0, 145.5, 160.4, 161.0, 162.6, and 180.3;  $\nu_{\text{max}} / \text{cm}^{-1}$  3249 (OH) and 1625 (C=O);  $\lambda_{\text{max}}(\text{CHCl}_3)$  nm (log  $\epsilon$ ) 276 (4.04), 292 (4.06), 300 (4.09), 321 (4.16), 339 (4.32, sh), 350 (4.38), 391 (3.80, sh), 396 (3.78, sh), 410 (3.68, sh), 428 (3.44, sh), 510 (3.20), 533 (3.18), and 581 (2.91, sh);  $\lambda_{\text{max}}(\text{MeOH})$  nm (log  $\epsilon$ ) 259 (3.77), 265 (3.82), 285 (3.95, sh), 297 (4.05, sh), 320 (4.23), 360 (4.29), 409 (3.65), 506 (3.79), and 531 (3.78);  $\lambda_{\text{max}}(\text{MeOH} + 1 \text{ drop of aq. NaOH})$  nm (log  $\epsilon$ ) 259 (3.77), 265 (3.82), 285 (3.94, sh), 298 (4.05, sh), 321 (4.25), 359 (4.28), 424 (3.65), 508 (3.79), and 532 (3.78). *Anal.* Calcd for  $\text{C}_{14}\text{H}_9\text{NO}_2 \cdot \text{EtOAc}$ : C, 69.44; H, 5.51; N, 4.50. Found: C, 69.41; H, 5.35; N, 4.24.

### Acetylation of 2f

A mixture of **2f** (0.0185 g, 0.083 mmol),  $\text{Ac}_2\text{O}$  (1 mL), and a drop of pyridine was stirred for 1 h at rt. To the mixture water was added, and the mixture was extracted with  $\text{CHCl}_3$ . The extract was dried over  $\text{Na}_2\text{SO}_4$ , and evaporated. The residue was chromatographed with EtOAc to give **2g** (0.0134 g, 61%) as violet powders.

**2g**: Violet needles (from MeOH- $\text{CH}_2\text{Cl}_2$ ), mp 188 °C;  $\delta_{\text{H}}$  2.44 (1H, s, Me), 7.10 (1H, d, *J* 12.2, H-4), 7.99 (1H, td, *J* 10.0 and 9.8, H-9), 8.05 (1H, dd, *J* 9.8 and 9.6, H-7), 8.10 (1H, s, H-1), 8.14 (1H, d, *J* 12.2, H-5), 8.17 (1H, t, *J* 9.8, H-8), 8.82 (1H, d, *J* 10.0, H-10), and 8.88 (1H, d, *J* 9.6, H-6);  $\delta_{\text{C}}$  20.8, 124.0, 126.3, 129.4, 139.5, 131.0, 132.9, 135.0, 139.2, 140.8, 144.6, 155.6, 159.2, 159.5, 168.8, and 180.0;  $\nu_{\text{max}} / \text{cm}^{-1}$  1754 and 1603 (C=O);  $\lambda_{\text{max}}(\text{CHCl}_3)$  nm (log  $\epsilon$ ) 259 (3.95), 288 (4.23, sh), 312 (4.36, sh), 327 (4.45), 340 (4.53), 360 (4.43), 384 (3.94, sh), 494 (3.82, sh), 503 (3.82), and 525 3.81, sh). *Anal.*

Calcd for C<sub>16</sub>H<sub>11</sub>NO<sub>3</sub>: C, 72.45; H, 4.18; N, 5.28. Found: C, 72.41; H, 4.33; N, 5.24.

### Reaction of 11-azacyclohept[*a*]azulen-3(3*H*)-ones (**2a**, **2b**, **2f**) with diethyl malonate

A mixture of **2a** (0.0872 g, 0.42 mmol) and diethyl malonate (0.35 mL, 2.3 mmol) in Ac<sub>2</sub>O (10 mL) was refluxed for 1.5 h. To the mixture was added water, then the mixture was extracted with CHCl<sub>3</sub>. The extract was dried over Na<sub>2</sub>SO<sub>4</sub>, and evaporated. The residue was chromatographed with EtOAc to give **4a** (0.318g, 24%) as red powders.

**4a**: Red powders (from CH<sub>2</sub>Cl<sub>2</sub>), mp 137-138 °C; δ<sub>H</sub> 1.45 (3H, t, *J* 7.1, Me), 4.43 (2H, q, *J* 7.1, OCH<sub>2</sub>), 7.07 (1H, dd, *J* 12.2 and 2.5, H-8), 7.22 (1H, dd, *J* 12.4 and 2.5, H-10), 7.51 (1H, dd, *J* 10.4 and 10.3, H-5), 7.73 (1H, dd, *J* 12.2, H-7), 7.81 (1H, dd, *J* 11.0 and 10.3, H-4), 8.04 (1H, d, *J* 10.4, H-6), 8.48 (1H, d, *J* 12.4, H-11), and 8.71 (1H, d, *J* 11.0, H-3); δ<sub>C</sub> 14.5, 60.4, 125.0, 126.5, 128.6, 129.4, 130.3, 130.9, 131.4, 135.2, 136.7, 136.8, 139.0, 140.7, 141.8, 149.6, 162.3, 163.4 and 187.3; ν<sub>max</sub> / cm<sup>-1</sup> 1738, 1674, 1619 (C=O), and 1588 (C=C); λ<sub>max</sub>(CHCl<sub>3</sub>) nm (log ε) 296 (4.32, sh), 334 (4.59), 341 (4.58), 376 (4.12, sh), 418 (3.23, sh), 518 (2.99, sh), 553 (3.04), and 604 (2.77, sh). *Anal.* Calcd for C<sub>19</sub>H<sub>13</sub>NO<sub>4</sub>: C, 71.47; H, 4.10; N, 4.39. Found: C, 71.53; H, 3.85; N, 4.21.

Similar treatment of **2b** and **2f** gave **4b** (39%) and **4f** (21%), respectively.

**4b**: Reddish brown powders (from cyclohexane), mp 223-224 °C; δ<sub>H</sub> 1.46 (3H, t, *J* 7.1, Me), 4.12 (3H, s, OMe), 4.45 (2H, q, *J* 7.1, OCH<sub>2</sub>), 7.24 (1H, d, *J* 12.4, H-8), 7.58 (1H, dd, *J* 10.3 and 9.8, H-5), 7.83 (1H, d, *J* 12.4, H-7), 7.85 (1H, dd, *J* 11.0 and 9.8, H-4), 8.11 (1H, d, *J* 10.3, H-6), 8.27 (1H, s, H-11), and 8.79 (1H, d, *J* 11.0, H-3); ν<sub>max</sub> / cm<sup>-1</sup> 1721, 1690, 1613 (C=O), and 1585 (C=C). *Anal.* Calcd for C<sub>20</sub>H<sub>15</sub>NO<sub>5</sub>: C, 68.76; H, 4.33; N, 4.01. Found: C, 69.02; H, 4.55; N, 3.86.

**4f**: Red powders (from CH<sub>2</sub>Cl<sub>2</sub>), mp 132-133 °C; δ<sub>H</sub> 1.45 (3H, t, *J* 7.1, Me), 4.43 (2H, q, *J* 7.1, OCH<sub>2</sub>), 7.25 (1H, d, *J* 12.4, H-8), 7.52 (1H, dd, *J* 10.4 and 9.6, H-5), 7.82 (1H, d, *J* 12.4, H-7), 7.83 (1H, dd, *J* 11.0 and 9.6, H-4), 8.05 (1H, d, *J* 10.4, H-6), 8.56 (1H, s, H-11), and 8.71 (1H, d, *J* 11.0, H-3) (OH was not observed); ν<sub>max</sub> / cm<sup>-1</sup> 3294 (OH), 1720, 1690, 1613 (C=O), and 1585 (C=C). *Anal.* Calcd for C<sub>19</sub>H<sub>13</sub>NO<sub>5</sub>: C, 68.06; H, 3.91; N, 4.18. Found: C, 67.72; H, 3.86; N, 4.28.

### Biological assay

HeLa S3 cells were obtained from AIST and used after cultivation. The cultivated HeLa S3 cells were cell counted and the culture fluid was prepared to the cell consistency of 2 × 10<sup>4</sup> cells/ml. The compounds added to the medium in DMSO solutions. To the aliquot of the culture fluid, which was incubated for 3 h at 37 °C, the test sample was added and then the culture fluid was incubated for 72 h.

To the culture fluid, MTT (3-[4,5-dimethylthiazol]-2-yl-2,5-diphenyltetrazolium bromide) solution was added, and incubated for 4 h. Then the sample was centrifuged at 3000 rpm for 10 min at 4 °C, and the solvent was evaporated. Then DMSO was added to the obtained mixture. The MTT-formazan was dissolved by plate-mixing and OD540 was measured. The rate of outlive determined to refer with un-dosed control. Dose-response curve was drawn up and IC<sub>50</sub> was pursued. Every experiment in the cycotoxic assay was replicated twice in order to define the IC values.

## REFERENCES AND NOTES

1. B. A. Hess, Jr. and L. J. Schaad, *J. Org. Chem.*, 1971, **36**, 3418.
2. T. Okazaki and K. K. Laali, *Org. Biomol. Chem.*, 2003, **1**, 3078.
3. A. Toyata and T. Nakajima, *Tetrahedron*, 1981, **37**, 2575.
4. M. Saito, T. Morita, and K. Takase, *Chem. Lett.*, 1974, 955.
5. M. Saito, T. Morita, and K. Takase, *Chem. Lett.*, 1975, 441.
6. M. Yasunami, T. Amemiya, and K. Takase, *Tetrahedron Lett.*, 1983, **24**, 69.
7. C. Jutz, H. G. Peuker, and W. Kosbahn, *Synthesis*, 1976, 673.
8. T. Toda, N. Shimazaki, T. Mukai, and C. Kabuto, *Tetrahedron Lett.*, 1980, **21**, 4001.
9. Z. Yoshida, M. Shibata, E. Ogino, and T. Sugimoto, *Tetrahedron Lett.*, 1984, **24**, 3343.
10. S. Kuroda, S. Hirooka, H. Iwaki, M. Ikeda, T. Nakao, M. Ogisu, M. Yasunami, and K. Takase, *Chem. Lett.*, 1986, 2039.
11. T. Takayasu and M. Nitta, *J. Chem. Soc., Perkin Trans. 1*, 1997, 3537.
12. For reviews see, N. Abe, 'Recent Research Developments in Organic and Bioorganic Chemistry' 2001, **4**, 14, Transworld Research Network; T. Nishiwaki and N. Abe, *Heterocycles*, 1981, **15**, 547; M. Kimura, *Yuki Gosei Kagaku Kyokai Shi*, 1981, **39**, 690.
13. N. Hirata, M. Yasunami, and K. Takase, *Tetrahedron Lett.*, 1975, 1849.
14. Y. Iino and M. Nitta, *J. Chem. Soc., Perkin Trans. 1*, 1994, 2579.
15. M. Nitta, Y. Iino, T. Sugiyama, and A. Toyota, *Tetrahedron Lett.*, 1993, **34**, 835.
16. M. Nitta, Y. Iino, T. Sugiyama, and A. Akaogi, *Tetrahedron Lett.*, 1993, **34**, 831.
17. K. Koizumi, C. Miyake, T. Ariyoshi, K. Umeda, N. Yamauchi, S. Tagashira, Y. Murakami, H. Fujii, and N. Abe, *Heterocycles*, 2007, **73**, 325.
18. M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, J. A. Montgomery, Jr., T. Vreven, K. N. Kudin, J. C. Burant, J. M. Millam, S. S. Iyengar, J. Tomasi, V. Barone, B. Mennucci, M. Cossi, G. Scalmani, N. Rege, G. A. Petersson, H. Nakatsuji, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, M. Klene, X. Li, J. E. Knox, H. P. Hratchian, J. B. Cross, C. Adamo, J. Jaramillo, R. Gomperts, R.

- E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, P. Y. Ayala, K. Morokuma, G. A. Voth, P. Salvador, J. J. Dennenberg, V. G. Zakrzewski, S. Dapprich, A. D. Daniels, M. C. Strain, O. Farkas, D. K. Malick, A. D. Rabuck, K. Raghavachari, J. B. Foresman, J. V. Ortiz, Q. Cui, A. G. Baboul, S. Clifford, I. Komaromi, R. L. Martin, D. J. Fox, T. Keith, M. A. Al-Laham, C. Y. Peng, A. Nanayakkara, M. Chllacombe, P. M. W. Gill, B. Johnson, W. Chen, M. W. Wong, C. Gonzalez, and J. A. Pople, *Gaussian 03*, Gaussian, Inc., Pittsburgh, PA (2003).
19. M. Oda, M. Funamizu, and Y. Kitahara, *Bull. Chem. Soc. Jpn.*, 1969, **42**, 2386.