

FORMATION OF 1,1'-BI-1H-PYRROLO[3,2-c]PYRIDINES BY THE DEOXYGENATION OF 4-NITRO-3-STYRYLPYRIDINE 1-OXIDES WITH TERVALENT PHOSPHORUS COMPOUNDS¹

Tarozaemon Nishiwaki,* Noritaka Abe, and Kenzo Moriyasu
Department of Chemistry, Faculty of Sciences, Yamaguchi University, Yamaguchi City 753, Japan


Abstract- The deoxygenation reaction of 4-nitro-3-styrylpyridine 1-oxides with triethyl phosphite or, more favourably, with hexaalkylphosphorous triamide afforded 1,1'-bi-1H-pyrrolo[3,2-c]pyridines instead of a monomeric product.

The N-N bonded dimers of pyrroles,² indoles,³ and carbazoles⁴ have been prepared by the oxidation reaction of the corresponding heterocycles. We wish to report that the deoxygenation reaction of 4-nitro-3-styrylpyridine 1-oxides with tervalent phosphorus compounds produces the N-N bonded dimers in moderate yields. This appears to be a rare instance in the deoxygenation reactions of nitro-arenes.⁵


Whereas 4-nitro-3-styrylpyridine 1-oxide⁶ (1a), when heated in a large excess of triethyl phosphite, gave 1-ethyl-2-phenyl-1H-pyrrolo[3,2-c]pyridine (2) [mp 198-199°, 29 % yield], heating of (1a) with 6 moles of triethyl phosphite in benzene for 7 days (the conditions reported to yield 2-phenyl-1H-pyrrolo[3,2-c]pyridine from (1a)⁷) afforded a compound C₂₆H₁₈N₄ [mp 266-268° (decomp.), 6 % yield].

¹H Nmr spectrum of this compound displays aromatic protons only whose chemical shifts and coupling constants are in keeping with those recorded for pyrrolo[3,2-c]pyridines⁸ and the spectral pattern did not change after the addition of deuterium oxide, indicating the absence of protons exchangeable with deuteriums. This is further supported by the absence of a ν(NH) absorption in its ir spectrum, which contains bands characteristic of pyrrolo[3,2-c]pyridine.⁹ From these observations the structure of 2,2'-diphenyl-1,1'-pyrrolo[3,2-c]pyridine (3a) was assigned.


The synthesis could be advantageously improved when (1a) was heated with 6 moles of hexamethyl- or hexaethyl-phosphorous triamide in *tert.*-butylbenzene or xylene at 130° in nitrogen for a few hours (a vigorous reaction took place at



(1)



(2)



(3)

(1), (3): a; R=H b; R=MeO c; R=Cl d; R=Me₂N e; R=Me

ca. 110° and the temperature of the mixture rose rapidly). By this procedure the following 2,2'-diaryl-1,1'-bi-1H-pyrrolo[3,2-c]pyridines¹⁰ were prepared: [(3a); 47 %]; [(3b); mp 278-279° (decomp.), 38 %]; [(3c); mp 290° (decomp.), 54 %]; [(3d); mp 280° (decomp.), 32 %]; [(3e); mp 272° (decomp.), 49 %].

Table 1. Spectral Data of 2,2'-Diaryl-1,1'-bi-1H-pyrrolo[3,2-c]pyridines (3a-e)

Compd.	λ_{max} . nm (log ϵ) (EtOH)	Ir Bands (cm ⁻¹) (Nujol)
(3a) ^a	242 (4.48), 248 (4.49), 304 (4.42)	1320, 910
(3b) ^b	253 (4.44), 260 (4.43), 309 (4.43)	1320, 910
(3c) ^c	244 (4.55), 251 (4.56), 309 (4.55)	1315, 910
(3d) ^c	225 (4.53), 279 (4.22), 334 (4.74)	1300, 910
(3e) ^c	250 (4.73), 306 (4.61)	1320, 910

^a δ [(CD₃)₂SO] 7.04 (s, 2H), 7.46 (m, 8H), 7.92 (m, 4H), 8.18 (d, J=6 Hz, 2H), 8.82 (s, 2H). ^b δ [(CD₃)₂SO] 3.80 (s, 6H), 6.90 (s, 2H), 7.07 (d, J=9 Hz, 4H), 7.34 (d, J=6 Hz, 2H), 7.84 (d, J=9 Hz, 4H), 8.15 (d, J=6 Hz, 2H), 8.78 (s, 2H). ^c Low solubilities of (3c-e) prevented nmr spectral determinations.

References and Notes

- Part 23 of Studies on Heterocyclic Chemistry. Part 22, T. Nishiwaki, E. Kawamura, N. Abe, and M. Iori, *Heterocycles*, in the press.
- T. Ajello and S. Giambone, *Gazzetta*, 1951, 81, 276.
- V. Dave, *Canad. J. Chem.*, 1972, 50, 3397.
- M. Sugita, *J. Chem. Soc. Japan*, 1967, 88, 659.
- J. I. G. Cadogan, *Synthesis*, 1969, 11.
- D. Jerchel and H. E. Heck, *Justus Liebigs Ann. Chem.*, 1958, 613, 171.
- M. H. Fisher, G. Schwartzkopf, and D. R. Hoff, *J. Med. Chem.*, 1972, 15, 1168.
- P. G. Riley and R. Robinson, *Canad. J. Chem.*, 1969, 47, 3257.
- R. E. Willette, *Adv. Heterocyclic Chem.*, 1968, 9, 27.
- Recrystallized from methanol. Satisfactory micro-analyses have been obtained.

Received, 24th March, 1980