

**REACTION OF 2-HYDRAZINO-1-AZAAZULENE WITH
DIPHENYLNITRONE: UNEXPECTED FORMATION OF
PHENYLIMINO-*p*-BENZOQUINONE-*N*-OXIDE**



**Noritaka Abe,^{*a} Haruhiko Matsuda,^a Hiroyuki Fujii^a and
Akikazu Kakehi^b**

^a Department of Chemistry, Faculty of Science, Yamaguchi University,
Yamaguchi 753-8512, Japan

^b Department of Chemistry and Material Engineering, Faculty of Engineering,
Shinshu University, Wakasato Nagano 380-8553, Japan

Abstract - Reaction of 2-hydrazino-1-azaazulene with diphenylnitrone gave benzaldehyde (1-azaazulen-2-yl)hydrazone (**2**) and phenylimino-*p*-benzoquinone-*N*-oxide (**3**). Reaction mechanism for the formation of **2** and **3** was discussed.

The chemistry of nitrone is extensively investigated because of its synthetically versatile use; an isoxazoline formation has been most worthy of remarks.¹ In the course of our investigation of azaazulene chemistry, we recently reported an interesting cyclization reaction where diphenylnitrone reacted with 2-alkylamino-1-azaazulenes and afforded novel pentacyclic heterocycles.² We also reported that 2-hydrazino-1-azaazulenes reacted with dimethyl acetylenedicarboxylate³ and diphenylcyclopropanone⁴ and gave some variety of cycloadducts and the hydrazones. From the expectation of the formation of new cycloadducts we expanded the investigation to the reaction of 2-hydrazino-1-azaazulenes with a nitrone. Some reports of phenylhydrazines with nitrones are appeared affording the corresponding hydrazones, but they were the reactions after hydrolysis of the nitrone.⁵ Reaction of 2-hydrazino-1-azaazulene (**1**) with diphenylnitrone in dry acetonitrile at rt for 3 days yielded benzaldehyde (1-azaazulen-2-yl)hydrazone (**2**) in 70% yield. When the reaction was performed under reflux for 3 days, a complex mixture was produced. From the mixture, two compounds, phenylimino-*p*-benzoquinone-*N*-oxide (**3**) (10.5% based on diphenylnitrone) and **2** (68%), were isolated. Compound (**2**) was deduced from the inspection of spectroscopic data as well as elemental analysis and independent synthesis; the reaction of **1** with benzaldehyde gave **2** in the 75% yield.



O ₁ —N ₁	1.29(1)	O ₁ —N ₁ —C ₁ —C ₂	172(1)°
O ₂ —C ₄	1.25(1)	O ₁ —N ₁ —C ₁ —C ₆	-6(1)°
N ₁ —C ₁	1.31(1)	O ₁ —N ₁ —C ₇ —C ₁₂	-59(1)°
C ₁ —C ₂	1.40(1)	O ₁ —N ₁ —C ₇ —C ₈	119.1(9)°
C ₁ —C ₆	1.44(1)	O ₂ —C ₄ —C ₅ —C ₆	179(1)°
C ₂ —C ₃	1.37(1)	C ₁ —N ₁ —C ₇ —C ₈	-61(1)°
C ₃ —C ₄	1.44(1)	C ₂ —C ₁ —N ₁ —C ₇	-8(2)°
C ₄ —C ₅	1.45(2)	C ₂ —C ₁ —C ₆ —C ₅	1(2)°
C ₅ —C ₆	1.34(1)	C ₃ —C ₂ —C ₁ —C ₆	2(2)°

Selective Bond Lengths (Å) and Torsional Angles

Figure 1. ORTEP drawing of **3** with thermal ellipsoid plot (50% probability)

The structure of **3** was confirmed by X-Ray structure analysis by direct method and an ORTEP drawing is shown in Figure 1.⁶ Compound (**3**) was first synthesized by Wieland and Roth,⁷ and attracted attentions; several investigations were reported about the synthesis, physical properties and reactions.^{8–11} The X-Ray structure analytical investigation of **3** itself has not been successful so far, only the report insufficient analysis of (*p*-methoxyphenyl)imino-*p*-benzoquinone-*N*-oxide was seen.¹²

Reaction of **1** with α -(*p*-methoxyphenyl)-*N*-phenylnitronone gave similar result as for **1**, and **3** (12%) and *p*-methoxybenzaldehyde (1-azaazulen-2-yl)hydrazone (**4**) (67%) were obtained.

A plausible mechanism is shown in the Scheme 1. The hydrazine (**1**) reacts with diphenylnitronone to give an adduct (**A**). Reaction of intermediate (**A**) with a second molar of diphenylnitronone causes cleavage of **A** and would give benzaldehyde (1-azaazulen-2-yl)hydrazone (**2**) and nitrosobenzene. The nitrosobenzene then reacts with **A** to give rise to **2** and the hydroxylamine (**B**). It is known that hydroxylamine underwent disproportionation and gave nitronone and amine,^{7,13} e.g., *N*-ethyl-*N*-phenylhydroxylamine gave α -methyl-*N*-phenylnitronone and *N*-ethylaniline.¹³ Moreover, it is known that the reaction of nitrosobenzene with *p*-benzoquinone gave dinitronone derivative.¹⁴ Therefore it is considered that disproportionation of **B** would give *p*-diiminoquinon-*N*-oxide (**C**). Addition of water to **C** gives **D**, and successive elimination of hydroxyamine furnishes **3**. Similar treatment of phenylhydrazine with diphenylnitronone did not give **3**. Therefore the results suggest a participation of 1-azaazulene moiety. Large electron-withdrawing character of the 1-azaazulene ring as shown in the resonance form **A'** would facilitate the reaction of **A** with nitrosobenzene.

EXPERIMENTAL

Melting points are uncorrected. ¹H NMR spectra (250 MHz) were recorded on a Hitachi R-250H spectrometer using deuteriochloroform as a solvent with tetramethylsilane as an internal standard; *J* values are recorded in Hz. Electronic spectra were taken with a Hitachi 220A spectrophotometer using ethanol as a solvent. IR spectra were recorded for KBr pellets on a Hitachi 270-50 infrared spectrophotometer. Kieselgel 60 was used for column chromatography.

Reaction of 2-hydrazino-1-azaazulene with diphenylnitronone

a) A solution of 2-hydrazino-1-azaazulene (**1**) (0.159 g, 1.00 mmol) and diphenylnitronone (0.591 g, 3.00 mmol) in dry acetonitrile (40 mL) was stirred at rt for 3 days and evaporated. The residue was chromatographed with chloroform to give benzaldehyde (1-azaazulen-2-yl)hydrazone (**2**) (0.174 g, 70%). Compound (**2**): orange prisms (from ethyl acetate), mp 223–224 °C; δ_{H} 7.10–7.30 (1H, br), 7.25–7.55 (7H, m), 7.71 (2H, d, *J* 7.0), and 7.80–8.20 (3H, m); ν_{max} / cm⁻¹ 1616 (C=N); λ_{max} nm (log ϵ) 252

(4.38), 288 (4.25), 340 (4.39), 410 (4.24), and 450 (4.13); m/z (rel. intensity) 248 ($M^+ + 1$, 24), 171 (81), 144 (90), and 90 (100); *Anal.* Calcd for $C_{16}H_{13}N_3$: C, 77.71; H, 5.30; N, 16.99. Found: C, 77.57; H, 5.34; N, 16.98.

b) A solution of 2-hydrazino-1-azaazulene (**1**) (0.477 g, 3.00 mmol) and diphenylnitrone (1.773 g, 9.00 mmol) in dry acetonitrile (60 mL) was refluxed for 3 days and evaporated. Chromatography of the residue with chloroform gave phenylimino-*p*-benzoquinone-*N*-oxide (**3**) (0.188 g, 10.5%) and **2** (0.504 g, 68%), successively. Compound (**3**): orange needles (from hexane–dichloromethane), mp 140–142 °C (lit.,² 140 °C); δ_H 6.26 (1H, dd, J 10.4 and 1.8), 6.67 (1H, dd, J 10.4 and 1.8), 7.19 (1H, dd, J 10.3 and 3.1), 7.45–7.60 (5H, m), and 8.05 (1H, dd, J 10.4 and 3.1); ν_{max} / cm^{-1} 1622 (C=O); λ_{max} nm (log ϵ) 261 (2.35), and 387 (3.03); m/z (rel. intensity) 200 ($M^+ + 1$, 15), 199 (M^+ , 15), and 92 (100); *Anal.* Calcd for $C_{12}H_9NO_2$: C, 72.35; H, 4.55; N, 7.03. Found: C, 72.09; H, 4.64; N, 7.05.

Synthesis of benzaldehyde (1-azaazulen-2-yl)hydrazone (**2**)

A solution of **1** (3.98 g, 25.0 mmol) and benzaldehyde (2.65 g, 25.7 mmol) in ethanol (100 mL) was refluxed for 1 h and evaporated. The residue was chromatographed with chloroform to give **2** (4.66 g, 75%).

Reaction of 2-hydrazino-1-azaazulene with α -(*p*-methoxyphenyl)-*N*-phenylnitrone

A solution of 2-hydrazino-1-azaazulene (**1**) (0.477 g, 3.00 mmol) and α -(*p*-methoxyphenyl)-*N*-phenylnitrone (1.704 g, 7.50 mmol) in acetonitrile (50 mL) was refluxed for 3 days and evaporated. The residue was chromatographed with chloroform to give *p*-methoxybenzaldehyde (1-azaazulen-2-yl)hydrazone (**4**) (0.540 g, 67%) and phenylimino-*p*-benzoquinone-*N*-oxide (**3**) (0.180 g, 12%).

Compound (**4**): orange prisms (from ethylacetate–hexane), mp 192–194 °C; δ_H 3.86 (3H, s), 6.94 (2H, d, J 8.7), 7.10–7.20 (1H, br), 7.30–7.42 (1H, br), 7.42–7.55 (1H, br), 7.68 (2H, d, J 8.7), 7.85–8.05 (2H, br m); ν_{max} / cm^{-1} 2830 (OCH₃) and 1614 (C=N); λ_{max} nm (log ϵ) 258 (4.33), 266 (4.42), 296 (4.37), 343 (4.47), 417 (4.36), and 452 (4.27); *Anal.* Calcd for $C_{17}H_{15}N_3O$: C, 73.63; H, 5.45; N, 15.15. Found: C, 73.72; H, 5.38; N, 15.33.

X-Ray structure determinations

Crystal data for 3: brown prism, $C_{12}H_9NO_2$, $M=199.21$, monoclinic, space group $P2_1/a$, $a=7.510(3)$, $b=12.6355(9)$, $c=10.302(2)$ Å, $\beta=101.08(2)^\circ$, $V=959.4(4)$ Å³, $Z=4$, $D_{calcd}=1.379$ g/cm³, crystal dimensions 0.24 x 0.88 x 1.00 mm. Data were measured on a Rigaku AFC 5S radiation diffractometer with graphite-monochromated Mo-K α radiation. A total 2476 reflections (2309 unique) were collected

using the ω - 2θ scan technique to a maximum 2θ value of 55.0° . The structure was solved by direct methods and refined by a full-matrix least-squares method using TEXAN structure analysis software,¹⁵ using 150 variables and 709 observed reflections ($I > 3\sigma(I)$). The non-hydrogen atoms were refined anisotropically. The weighting scheme $\omega = 4Fo^2/\sigma^2(Fo^2)$ gave satisfactory agreement analyses. The final R and Rw values were 0.093 and 0.115. The maximum peak and the minimum peak in final difference map were $0.48 \text{ e}/\text{\AA}^3$ and $-0.40 \text{ e}/\text{\AA}^3$.

REFERENCES

1. L. I. Smith, *Chem. Rev.*, 1938, **23**, 193; J. Hamer and A. Macaluso, *Chem. Rev.*, 1964, **64**, 473; J. J. Tufariello, *1,3-Dipolar Cycloaddition Chemistry*, ed. by A. Padwa, John Wiley & Sons, New York, 1984, Vol. 2, chap. 9; A. Padwa and A. M. Schoffstall, *Advances in Cycloaddition*, ed. by D. P. Curran, JAI Press Inc., Greenwich, London, 1990, Vol. 2, p. 2.
2. N. Abe, H. Matsuda, Y. Sugihara, and A. Kakehi, *J. Chem. Soc., Perkin Trans. 1*, 1997, 993.
3. N. Abe and Y. Emoto, *Bull. Chem. Soc. Jpn.*, 1988, **61**, 1440.
4. N. Abe and A. Kakehi, *Bull. Chem. Soc. Jpn.*, 1994, **67**, 2487.
5. A. Giner-Soreolla, I. Zimmermann, and A. Bendich, *J. Am. Chem. Soc.*, 1959, **81**, 2515.
6. C. K. Johnson, "ORTEP II, Report ORNL-5138," Oak Ridge National Laboratory, Oak Ridge, Tennessee, 1976.
7. H. Wieland and K. Roth, *Ber.*, 1920, **53B**, 210.
8. C. J. Pedersen, *J. Am. Chem. Soc.*, 1957, **79**, 5014.
9. G. R. Chalfont, D. H. Hey, K. S. Y. Liang, and M. J. Perkins, *J. Chem. Soc. (B)*, 1971, 233.
10. A. R. Forrester, M. M. Ogilvy, and R. H. Thomson, *J. Chem. Soc., Chem. Commun.*, 1972, 483; A. R. Forrester, M. M. Ogilvy, and R. H. Thomson, *J. Chem. Soc., Perkin Trans. 1*, 1982, 2023; P. J. Baldry, A. R. Forrester, M. M. Ogilvy, and R. H. Thomson, *J. Chem. Soc., Perkin Trans. 1*, 1982, 2027; P. J. Baldry, A. R. Forrester, M. M. Ogilvy, and R. H. Thomson, *J. Chem. Soc., Perkin Trans. 1*, 1982, 2035.
11. J. A. Damavandy and R. A. Y. Jones, *J. Chem. Soc., Perkin Trans. 1*, 1981, 712.
12. C. Romers and B. Hesper, *Acta Cryst.*, 1966, **20**, 162.
13. G. Renner, *Z. Anal. Chem.*, 1963, **193**, 92.
14. W. Guendel and R. Pummerer, *Ann.*, 1937, **529**, 11.
15. TEXAN TEXRAY, Structure Analysis Package, Molecular Structure Corporation, 1985.