

CYCLOADDITION OF 1-METHYL-1-AZAAZULEN-2(1H)-ONE WITH  
REACTIVE ACETYLENES

Noritaka Abe\* and Toshiyuki Takehiro







Department of chemistry, Faculty of Sciences, Yamaguchi  
University, Yamaguchi 753, Japan

Abstract—Reactions of 1-methyl-1-azaazulen-2(1H)-one with reactive acetylenes in refluxing acetonitrile gave 1,2-disubstituted azulene (2), 2-methylcyclopent[de]isoquinolin-3(2H)-one (3), 1-methylcyclopent[cd]azulen-2(1H)-ones (4 and 5), and 3-substituted 1-methyl-1-azaazulen-2(1H)-ones (6 and 7). When the reaction was carried out in refluxing t-butylbenzene, the compound 2 and 6,8a-etheno-1-methyl-1-azaazulen-2(1H)-one (8) were obtained as major products.



We have reported the cycloadditions of 1-azaazulenes with reactive acetylenes, which proceeded via dipolar intermediates.<sup>1</sup> As a continuation of our work, we advanced the reaction to 1-methyl-1-azaazulen-2(1H)-one (1) and found that 1 and reactive acetylenes underwent interesting cycloaddition reactions and rearrangement.

Treatment of 1 with dimethyl acylenedicarboxylate (DMAD) in refluxing acetonitrile for 120 h gave dimethyl 1,2-azulenedicarboxylate (2a)<sup>2</sup> (9%), dimethyl 2-methyl-2,3-dihydro-3-oxocyclopent[de]isoquinoline-4,5-dicarboxylate (3a)<sup>3</sup> (11%), dimethyl 1-methyl-1,2,2a,3-tetrahydro-2-oxo-1-azacyclopent[cd]azulene-3,4-dicarboxylate (4a)<sup>4</sup> (11%), dimethyl 1-methyl-1,2-dihydro-2-oxo-1-azacyclopent[cd]azulene-3,4-dicarboxylate (5a)<sup>5</sup> (7%), dimethyl 1-methyl-1,2-dihydro-2-oxo-1-azaazulene-3-maleate (6a)<sup>6</sup> (17%), and dimethyl 1-methyl-1,2-dihydro-2-oxo-1-azaazulene-3-fumalate (7a)<sup>7</sup> (21%). These structures were determined on the basis of the spectroscopic data as well as elemental analyses. Compound 3a was a stable condensed isoquinolone derivative. In its <sup>1</sup>H-nmr spectrum, a signal of 1H singlet assignable to H-1 proton is seen at  $\delta$  8.73, which is a reasonable value as 3-isoquinolone.<sup>8</sup> In the <sup>1</sup>H-nmr spectrum of 4a, AB doublets are seen at  $\delta$  3.10 and 4.32 (J=9.2 Hz), and in the <sup>13</sup>C-nmr spectrum, two methine signals at  $\delta$  43.14 and

a: E = CO<sub>2</sub>Me , b: E = COPh



7a,b: (E)-isomer



Scheme 1

50.09. Dehydrogenation of 4a by tetrachloro-o-benzoquinone gave 5a in a 47% yield. The results associate with the structures.

When the above reaction was carried out in refluxing t-butylbenzene for 3 h, compounds 2a (21%), 3a (5%), 5a (0.8%), and dimethyl 6,8a-etheno-1-methyl-1,2-dihydro-2-oxo-1-azaazulene-7,8-dicarboxylate (8a)<sup>9</sup> (49%), were isolated. In the <sup>13</sup>C-nmr spectrum of 8a, signals assignable to sp<sup>3</sup> carbons are seen at  $\delta$  38.70 (d, C-6) and 71.55 (s, C-8a), and in its <sup>1</sup>H-nmr spectrum, methine proton is observed at  $\delta$  4.46 (ddd, J=8.5, 6.7, and 1.2 Hz, H-6). Further evidence supporting the structure of 8a is provided by its mass spectrum which displays an intense peak at m/z 159 (98%) associated with the loss of DMAD.

In a similar treatment, a reaction of 1 with dibenzoylacetylene in refluxing acetonitrile gave 2b (9%), 3b (3%), 4b (50%), 5b (6%), and 7b (6%), and in refluxing t-butylbenzene 2b (60%), 3b (19%), 5b (1.6%), and 8b (15%).<sup>10</sup>

A plausible mechanism is shown in Scheme 1. When DMAD attacks at C-3 position of 1, which is a reactive site towards electrophile,<sup>11</sup> a dipolar species A should be produced. A cyclization on C-8a position of A and a successive elimination of MeNCO furnishes 2 (path a). A cyclization on C-4 position of A gives B (path b), and a subsequent 1,5-hydrogen shift of B produces C (path c) and 4 (path d). Dehydrogenation of B and/or 4 affords 5. A tautomerization of C and a subsequent dehydrogenation lead to 3 (path c). A hydrogen shift of A gave Michael addition products 6 and 7 (path e). On the reaction in refluxing t-butylbenzene a Diels-Alder reaction occurs on seven-membered ring and 8 is obtained, in addition to the favorable cycloaddition-cycloreversion product 2. Similar periselectivity dependent on temperature is known on the reaction of 5-azaazulene with DMAD.<sup>12</sup>

We thank Dr. Masafumi Yasunami (Tohoku University) and Dr. Akira Mori (Kyushu University) for the measurements of elemental analyses and mass spectra.

#### REFERENCES

1. N. Abe, Y. Tanaka, and T. Nishiwaki, *J. Chem. Soc., Perkin Trans. 1*, 1978, 429; N. Abe and T. Nishiwaki, *Bull. Chem. Soc. Jpn.*, 1981, 54, 1277.
2. mp 53-54°C (lit. 2a, oil; 2b, mp 45°C), MS m/z 244 (M<sup>+</sup>, 81%), 213 (100), <sup>1</sup>H-nmr (250 MHz)  $\delta$  =3.96 (s, OMe), 3.99 (s, OMe), 7.48 (t, J=9.8 Hz, H-5), 7.48 (s, H-3), 7.57 (t, J=9.8 Hz, H-7), 7.87 (t, J=9.8 Hz, H-6), 8.50 (d, J=9.8 Hz, H-4), 9.44 (d, J=9.8 Hz, H-8); 2a. W. von E. Doering and D. W. Wiely, *Tetrahedron*, 1960, 11, 183; 2b. I. Kawamoto, Y. Sugimura, N. Soma, and Y. Kishida, *Chem. Lett.*, 1972, 931.

3. mp 168-169°C, MS m/z 299 (M<sup>+</sup>, 100%), <sup>1</sup>H-nmr (250 MHz) δ =3.40 (s, NMe), 3.99 (s, OMe), 4.11 (s, OMe), 6.95 (dd, J=6.1 and 1.2 Hz, H-6), 7.45-7.60 (m, H-7 and 8), and 8.73 (s, H-1), <sup>13</sup>C-nmr (62.87 MHz) δ =26.36 (q), 52.84 (q), 53.08 (q), 106.85 (d), 120.59 (d), 123.86 (s), 126.08 (s), 127.81 (s), 127.99 (s), 130.21 (d), 130.50 (s), 134.04 (d), 139.85 (s), 165.30 (s), 165.38 (s), and 166.64 (s).
4. mp 151-152°C, MS m/z 301 (M<sup>+</sup>, 100%), <sup>1</sup>H-nmr (250 MHz) δ =3.08 (s, NMe), 3.10 (d, J=9.2 Hz, H-2a), 3.86 (s, OMe), 3.88 (s, OMe), 4.32 (d, J=9.2 Hz, H=3), 5.66 (d, J=6.1 Hz, H-8), 6.40-6.50 (m, H-5 and 6), and 6.65-6.75 (m, H-7), <sup>13</sup>C-nmr (62.87 MHz) δ =27.14 (q), 43.14 (d), 50.09 (d), 52.33 (q), 52.55 (q), 98.33 (d), 122.19 (d), 124.69 (d), 124.75 (s), 130.83 (s), 131.29 (d), 134.43 (s), 134.68 (s), 162.86 (s), 164.40 (s), and 170.73 (s).
5. mp 190-191°C, MS m/z 299 (M<sup>+</sup>, 35%), and 126 (100), <sup>1</sup>H-nmr (250 MHz) δ =3.52 (s, NMe), 3.94 (s, OMe), 4.05 (s, OMe), 7.48 (d, J=9.8 Hz, H-8), 7.72 (t, J=9.8 Hz, H-6), 8.15 (t, J=9.8 Hz, H-7), and 9.16 (d, J=9.8 Hz, H-5), <sup>13</sup>C-nmr (62.87 MHz) δ =26.55 (q), 51.67 (q), 52.72 (q), 113.74 (d), 114.03 (s), 115.22 (s), 128.29 (d), 135.17 (s), 137.11 (s), 138.93 (s), 140.87 (d), 141.40 (d), 149.96 (s), 162.79 (s), 164.23 (s), and 180.28 (s).
6. mp 123.5-124°C, <sup>1</sup>H-nmr (250 MHz) δ =3.58 (s, NMe), 3.79 (s, OMe), 3.96 (s, OMe), 7.00 (s, H-vinyl), 7.05-7.40 (m, H-5, 6, 7, and 8), 7.76 (d, J=11.0 Hz, H-4).
7. mp 150-151°C, <sup>1</sup>H-nmr (250 MHz) δ =3.58 (s, NMe), 3.68 (s, OMe), 3.81 (s, OMe), 6.90-7.30 (m, H-4, 5, 6, 7, and 8), and 7.15 (s, H-vinyl).
8. H. Fukumi and H. Kurihara, Heterocycles, 1978, 9, 1197.
9. mp 156-158°C, MS m/z 301 (M<sup>+</sup>, 100%), and 159 (98), <sup>1</sup>H-nmr (250 MHz) δ =3.05 (s, NMe), 3.77 (s, OMe), 3.78 (s, OMe), 4.46 (ddd, J=8.5, 6.7, and 1.2 Hz, H-6), 5.95 (s, H-3), 6.06 (d, J=10.4 Hz, H-4), 6.21 (dd, J=7.9 and 1.2 Hz, H-9), 6.41 (dd, J=10.4 and 8.5 Hz, H-5), and 6.66 (dd, J=7.9 and 6.7 Hz, H-10), <sup>13</sup>C-nmr (62.87 MHz) δ =26.47 (q), 38.70 (d), 52.64 (q, x 2), 71.55 (s), 120.57 (d), 121.16 (d), 129.47 (d), 132.62 (d), 133.26 (d), 134.23 (s), 145.64 (s), 147.03 (s), 163.07 (s), 164.95 (s), and 171.28 (s).
10. The spectral data of the products other than recorded in this text showed pertinent figures, and the full data will be reported elsewhere.
11. T. Toda, S. Ryu, Y. Hagiwara, and T. Nozoe, Bull. Chem. Soc. Jpn., 1975, 48, 82; H. Nakao, N. Soma, Y. Sato, and G. Sunagawa, Chem. Pharm. Bull., 1965, 13, 473.
12. K. Hafner and J. Haring, Erdoel Kohle, Erdgas, Petrochem., 1979, 32, 136.

Received, 18th February, 1987